arXiv:hep-th/0103151v1 19 Mar 2001

Bouncing Branes

Emil M. Prodanov[*]

National Centre for Scientific Research "Demokritos", Athens, Greece

Abstract

Two classical scalar fields are minimally coupled to gravity in the Kachru–Shulz–Silverstein scenario with a rolling fifth radius. A Tolman wormhole solution is found for a $\mathbb{R} \times S^3$ brane with Lorentz metric and for a $\mathbb{R} \times AdS_3$ brane with positive definite metric.

PACS numbers: 04.50.+h, 11.27.+d, 98.80.Cq.
Keywords: Wormhole, Tolman, Extra Dimension, Localization of Gravity, Kachru–Shulz–Silverstein.

---

[*] e-mail: prodanov@maths.tcd.ie

1 Introduction

Two classical scalar fields are minimally coupled to gravity in the Kachru–Shulz–Silverstein scenario [1] with a rolling fifth radius. One of these fields is time-dependent only, while the other is the dilaton. We consider the most general four-dimensional homogeneous and isotropic Robertson–Walker metric [2] (with a time-dependent scale factor), naturally generalized to a five-dimensional Randall–Sundrum [3] context. We find a Tolman wormhole solution [4] for a $\mathbb{R}\times\mathrm{S}^3$ brane with Lorentz metric and for $\mathbb{R}\times\mathrm{AdS}_3$ brane with positive definite metric. The solution represents a collapsing Universe which starts expansion just before encountering a big crunch singularity. The Universe reaches a moment of minimal spatial volume. This minimum volume edgeless achronal spacelike hypersurface is called a *bounce* [5]. The wormhole is time-dependent and the bounce involves the entire Universe.

Our analysis will be along the lines of the one in [6].

2 The Model

The action for gravity coupled to two scalars is:

$$S \;=\int d^4x\, dr\;\left(\mathcal{L}^{(5)}_{MATTER}\;+\;\mathcal{L}^{(5)}_{GRAVITY}\right), \tag{1}$$

where:

$$\mathcal{L}^{(5)}_{MATTER} = -\frac{1}{2}\sqrt{|g^{(5)}|}\nabla^\mu\phi\nabla^\nu\phi\, g^{(5)}_{\mu\nu} - \sqrt{|g^{(5)}|}U(\phi) - \sqrt{|g^{(4)}|}V(\phi)\delta(r)\,, \tag{2}$$

$$\mathcal{L}^{(5)}_{GRAVITY} = \frac{1}{\kappa^2}\sqrt{|g^{(5)}|}\;R. \tag{3}$$

Here, $g^{(4)}_{\mu\nu}$ is the pull-back of the five-dimensional metric $g^{(5)}_{\mu\nu}$ to the domain wall taken to be at $r=0$. The wall is represented by a delta-function source with coefficient $V(\phi)$, parametrising its tension. The scalar $\phi(t,r)$ is:

$$\phi(t,r) \;=\; \psi(t) \;+\; \chi(r)\,. \tag{4}$$

We make the ansatz that both the potentials U and V are of Liouville type:

$$U(\phi) \;=\; U_0 e^{\alpha\phi}\,, \tag{5}$$

$$V(\phi) \;=\; V_0 e^{\beta\phi} \tag{6}$$

and that the metric is:

$$ds^2 = se^{-A(r)}dt^2 + \frac{e^{-A(r)}g(t)}{\left[1+\frac{\epsilon}{4}(x^2+y^2+z^2)\right]^2}\,(dx^2+dy^2+dz^2) + qf(t)dr^2\,. \tag{7}$$

This is a natural generalisation of the most general four-dimensional homegeneous and isotropic Robertson–Walker metric [2] to a five-dimensional Randall–Sundrum context [3]. The scale factor $g(t)$ is a strictly positive function (we are working with a mostly-plus metric) and the function $f(t)$ is strictly positive as well (the metric is never degenerate).The factors s and q are signs ($s^2 = q^2 = 1$). The curvature parameter is $\epsilon = +1$ (for spherical spatial three-sections) or $\epsilon = -1$ (for hyperbolical spatial three-sections).
The energy–momentum tensor for the scalar fields is:

$$\begin{aligned} T_{\mu\nu} \;=\; & \frac{1}{2}\,\nabla_\mu\phi\,\nabla_\nu\phi \,-\, \frac{1}{2}\,g^{(5)}_{\mu\nu}\left[\frac{1}{2}\,\nabla_\alpha\phi\,\nabla_\beta\phi\,g^{(5)\alpha\beta} \,+\, U(\phi)\right] \\ & -\frac{1}{2}\,\frac{\sqrt{|g^{(4)}|}}{\sqrt{|g^{(5)}|}}\,V(\phi)\,\delta(r)\,g^{(4)}_{ab}\,\delta^a_\mu\,\delta^b_\nu\,. \end{aligned} \tag{8}$$

Einstein's equations $G_{\mu\nu} = \kappa^2 T_{\mu\nu}$, equivalently written in terms of the Ricci tensor as $R_{\mu\nu} = \kappa^2(T_{\mu\nu} - \frac{1}{3}g^{(5)}_{\mu\nu}T^{\alpha}{}_{\alpha})$, are:

$$\begin{aligned} & -\frac{s}{q}\frac{1}{f}e^{-A}A'^2 + \frac{s}{2q}\frac{1}{f}e^{-A}A'' + \frac{1}{4}\frac{\dot{f}^2}{f^2} - \frac{1}{2}\frac{\ddot{f}}{f} + \frac{3}{4}\frac{\dot{g}^2}{g^2} - \frac{3}{2}\frac{\ddot{g}}{g} \\ & \qquad = \frac{\kappa^2}{2}\dot{\psi}^2 + \frac{\kappa^2}{3}se^{-A}U + \frac{\kappa^2}{6}\frac{s}{f^{1/2}}e^{-A}V\delta(r)\,, \end{aligned} \tag{9}$$

$$\begin{aligned} & 2\epsilon - \frac{1}{q}\frac{g}{f}e^{-A}A'^2 + \frac{1}{2q}\frac{g}{f}e^{-A}A'' - \frac{1}{4s}\frac{\dot{f}}{f}\dot{g} - \frac{1}{4s}\frac{\dot{g}^2}{g} - \frac{1}{2s}\ddot{g} \\ & \qquad = \frac{\kappa^2}{3}e^{-A}gU + \frac{\kappa^2}{6}e^{-A}\frac{g}{f^{1/2}}V\delta(r)\,, \end{aligned} \tag{10}$$

$$\begin{aligned} & -A'^2 + 2A'' + \frac{q}{4s}\frac{\dot{f}^2}{f}e^{A} - \frac{q}{2s}\ddot{f}e^{A} - \frac{3q}{4s}\dot{f}\frac{\dot{g}}{g}e^{A} \\ & \qquad = \frac{\kappa^2}{2}\chi'^2 + \frac{\kappa^2}{3}qfU + \frac{2\kappa^2}{3}qf^{1/2}V\delta(r)\,, \end{aligned} \tag{11}$$

$$-\frac{3}{4}A'\frac{\dot{f}}{f} \;=\; \frac{\kappa^2}{2}\dot{\psi}\chi'\,. \tag{12}$$

The tr-equation, (12), implies:

$$\kappa\dot{\psi}(t) \;=\; -\sqrt{3}\;\frac{\dot{f}(t)}{f(t)}\,, \tag{13}$$

$$\kappa\chi'(r) \;=\; \frac{\sqrt{3}}{2}\;A'(r)\,. \tag{14}$$

When $\beta = \frac{\kappa}{2\sqrt{3}}$, the potential $V(\phi)$ can be written in the form:

$$V\Big(\phi(t,r)\Big) \;=\; \frac{1}{\kappa^2}\frac{1}{qf^{1/2}}W\,, \tag{15}$$

where W is a constant.
Let us assume that the potential $U(\phi)$ can be written as a function of t and r in the separable form:

$$U\Big(\phi(t,r)\Big) \;=\; \frac{1}{\kappa^2}\left[\frac{1}{qf}U_1(r) + e^A U_2(t)\right]. \tag{16}$$

At the end we will recast the potential U back into the original exponential form (5).
Einstein's equations then reduce to:

$$\frac{1}{4}\frac{\dot f^2}{f^2} - \frac{1}{2}\frac{\ddot f}{f} + \frac{1}{4}\frac{\dot f}{f}\frac{\dot g}{g} + \frac{\dot g^2}{g^2} - \frac{\ddot g}{g} - \frac{2\epsilon s}{g} - \frac{3}{2}\frac{\dot f^2}{f^2} \;=\; 0\,, \tag{17}$$

$$-\frac{1}{4s}\frac{\dot f}{f}\frac{\dot g}{g} - \frac{1}{4s}\frac{\dot g^2}{g^2} - \frac{1}{2s}\frac{\ddot g}{g} + \frac{2\epsilon}{g} - \frac{1}{3}U_2 \;=\; \frac{C}{qf}\,, \tag{18}$$

$$\frac{1}{2}\frac{\dot f^2}{f^2} - \frac{3}{2}\frac{\dot f}{f}\frac{\dot g}{g} - \frac{\ddot f}{f} - \frac{2s}{3}U_2 \;=\; \frac{2sD}{qf}\,, \tag{19}$$

$$A'^2 - \frac{1}{2}A'' + \frac{1}{3}U_1 + \frac{1}{6}W\delta(r) \;=\; Ce^A\,, \tag{20}$$

$$\frac{11}{8}A'^2 - 2A'' + \frac{1}{3}U_1 + \frac{2}{3}W\delta(r) \;=\; De^A\,, \tag{21}$$

where C and D are separation constants.
We will be looking for a bounce solution in the form:

$$g(t) \;=\; f(t) \;=\; Bt^2 + h \;>\; 0\,, \tag{22}$$

where B is a positive constant, not equal to 1 (so that the pull-back of the metric to 4 dimensions is flat only asymptotically*), and h is another strictly positive constant.
Upon substitution of the solutions (22) into the Einstein's equations, (17) gives:

$$B \;=\; -\frac{2\epsilon s}{3}\,. \tag{23}$$

---

*The curvature of the brane is $R = \frac{6-6B}{Bt^2+h}$.

On the otherhand, B must be positive. Therefore $B = \frac{2}{3}$ and ϵ and s must have opposite signs. Thus the solution is either a brane with spherical three-sections and Lorentz metric or a brane with hyperbolical three-sections and positive definite metric. Clearly, these two solutions can also be related by a Wick rotation (time t is changed to it and the positive-curvature spacetime becomes a negative-curvature spacetime).
Einstein's equations (18) and (19) are consistent if we choose

$$U_2(t) \;=\; \sigma \frac{\dot{g}^2(t)}{g^2(t)}\,, \tag{24}$$

where σ is a constant.
The next Einstein's equation, (18), yields that the separation constant C is $\frac{8\epsilon q}{3}$ and that $\sigma = -\frac{3}{2s}$.
The remaining time-dependent Einstein's equation, (19), gives $D = \frac{C}{4} = \frac{2\epsilon q}{3}$.
The r-dependent Einstein's equations (20) and (21) yield:

$$U_1(r) \;=\; 10\epsilon q e^{A(r)} - \frac{21}{8}A'(r)^2 \tag{25}$$

and these two equations reduce to a single equation:

$$\frac{1}{8}A'^2 - \frac{1}{2}A'' + \frac{1}{6}W\delta(r) \;=\; -\frac{2\epsilon q}{3}e^A\,. \tag{26}$$

A solution of this equation is of KSS [1] type:

$$A(r) \;=\; \ln\frac{1}{(k|r|+1)^2}\;, \tag{27}$$

where k is a constant, such that $k^2 = \frac{4\epsilon q}{3}$. Therefore ϵ and q must have the same signs ($k^2 = 4/3$). The constant W in brane tension V is $-12k$.
The equation of motion for the scalar field:

$$\nabla^2\phi \;-\; \frac{\delta U(\phi)}{\delta\phi} \;-\; \frac{\sqrt{|g^{(4)}|}}{\sqrt{|g^{(5)}|}}\,\frac{\delta V(\phi)}{\delta\phi}\,\delta(r) \;=\; 0\,, \tag{28}$$

after integration over the fifth dimension in an infinitesimal interval, gives a jump condition acrross the brane:

$$A'(+0) - A'(-0) \;=\; -4k\,. \tag{29}$$

Let us now write the potential $U\Big(\phi(t,r)\Big) = \frac{1}{\kappa^2}\left[\frac{1}{qf}U_1(r) + e^A U_2(t)\right]$ back in the exponential form $U = U_0 e^{\alpha\phi}$. Substituting the solution (27) for $A(r)$ into (25) gives:

$$U_1(r) \;=\; -4\epsilon q e^{A(r)}\,. \tag{30}$$

Using this form of $U_1(r)$, together with (24) for $U_2(t)$ and the value of B, we easily find that:

$$U\Big(\phi(t,r)\Big) \;=\; -\frac{4\epsilon h}{\kappa^2}\,\frac{e^A}{g^2} \;=\; U_0 e^{\frac{2\kappa}{\sqrt{3}}\phi}\,. \tag{31}$$

For a realistic model, one has to choose h sufficiently small.

Acknowledgements

It is a pleasure to thank B. Dolan, V. Gueorguiev, G. Savvidy, S. Sen and especially C. Kennedy for many useful discussions. Comments from C. P. Burgess, S. Giddings, I. Novikov, A. Strominger and N. Tetradis were also very helpful. The financial support of the EU, grant: HPRN–CT–1999–00161, is gratefully acknowledged.

References

[1] S. Kachru, M. Schulz and E. Silverstein: *Self-tuning Flat Domain Walls in 5d Gravity and String Theory*. Phys. Rev. **D62** (2000) 045021, hep-th/0001206;
S. Kachru, M. Schulz and E. Silverstein: *Bounds on Curved Domain Walls in 5-D Gravity*. Phys. Rev. **D62** 085003 (2000), hep-th/0002121.

[2] Ch. W. Misner, K. S. Thorne and J. A. Wheeler: **Gravitation**. W. H. Freeman & Co. (1973).

[3] L. Randall and R. Sundrum: *A Large Mass Hierarchy from a Small Extra Dimension*. Phys. Rev. Lett. **83**, 3370–3373 (1999), hep-ph/9905221;
L. Randall and R. Sundrum: *An Alternative to Compactification*. Phys. Rev. Lett. **83**, 4690–4693 (1999), hep-th/9906064.

[4] A. Einstein: Berl. Ber. **235** (1931);
R. C. Tolman: Phys. Rev. **38**, 1758 (1931)

[5] D. Hochberg, C. Molina-Paris and M. Visser: *Tolman Wormholes Violate the Strong Energy Condition*. Phys. Rev. **D59**, 044011 (1999), gr-qc/9810029.

[6] C. Kennedy and E. M. Prodanov: *Standard Cosmology From Sigma Model*. Phys. Lett. **B488** 11–16 (2000), hep-th/0003299;
C. Kennedy and E. M. Prodanov: *Standard Cosmology on a Self-Tuning Domain Wall*. hep-th/0010202.